

Ethnobotanical, Ethnopharmacological, and Phytochemical Studies of *Myrtus communis* Linn: A Popular Herb in Unani System of Medicine

Journal of Evidence-Based
Complementary & Alternative Medicine
2017, Vol. 22(4) 1035-1043
© The Author(s) 2017
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/2156587217718958
journals.sagepub.com/home/cam



Mekonnen Sisay, MSc¹ and Tigist Gashaw, MSc¹

Abstract

Myrtus communis L (Myrtaceae) is one of the popular drugs being used in the Unani system of phytomedicine since ancient Greece period. From time immemorial, different parts of this plant and essential oil have been used for a variety of purposes such as cosmetics (hair fall control), flavoring of food and drinks as well as extensive therapeutic purposes. Ethnobotanical information revealed that *M communis* L has been a folkloric reputed for the treatment of several diseases like gastric ulcer, diarrhea, dysentery, cancer, rheumatism, hemorrhage, deep sinuses, leucorrhoea, hemorrhoid, inflammation, dyspepsia, anxiety, insomnia, diabetes, hypertension, pulmonary disorders, and skin diseases. Moreover, ethnopharmacological studies revealed that the plant is endowed with extensive pharmacological activities, including antimicrobial, antidiarrheal, antidiabetic, antispasmodic, vasodilator, antiulcer, antioxidant, anticancer, anxiolytic, sedative-hypnotic, and anti-inflammatory activities, among others. The plant has been known to contain phenolic acids, tannins, flavonoids, glycosides, and terpenes. The myrtle oil was also found to be rich in a variety of bioactive monoterpenes and sesquiterpenes with their derivatives. Most of these studies validate the aforementioned traditional claims of this medicinal plant. Further studies are needed to unravel other pharmacological activities of this plant in the long run.

Keywords

Myrtus communis L, ethnobotanical uses, ethnopharmacological studies, phytochemical studies

Received March 15, 2017. Accepted for publication June 5, 2017.

Botanical Source and Characteristics

Family Myrtaceae includes approximately 100 genera and 3000 species growing in temperate, tropical, and subtropical regions of the world. *Myrtus* is a small genus belonging to this family¹ (Table 1). The genus *Myrtus* L comprises 2 species, *Myrtus communis* L (common myrtle) growing wild all around the Mediterranean basin and *Myrtus nivellei* Batt (Saharan myrtle) found in central Sahara. It is an aromatic evergreen perennial shrub² (Figure 1).

Methods

Data were collected from several legitimate data bases and services such as PubMed, Medline, Scopus, and other database sources like Google Scholar using key terms “*Myrtus communis* L”, “ethnobotanical uses”, “ethnopharmacological studies”, “phytochemical studies” in each database. Relevant and related data were filtered properly if it was found appropriate to the topic of interest. Time frame was also adjusted to obtain up-to-date information

regarding *M communis* L and its role in Unani system of medicine, ethnobotanical, ethnopharmacological, and phytochemical aspects of it by emphasizing on original researches conducted and published from 2010 to 2017. Excluding the background information, around 73 references were included in this scientific review. The study was conducted from September 2016 to January 2017.

Ethnobotanical Uses

The common myrtle is one of the most important drugs being used in Unani system of medicine, which originated in Greece

¹ Haramaya University, Harar, Ethiopia

Corresponding Author:

Mekonnen Sisay, Department of Pharmacology, School of Pharmacy, College of Health and Medical Sciences, Haramaya University, Harar, PO Box 235, Ethiopia.

Email: mekonnensisay27@yahoo.com



Table 1. Ethnobotanical Sources and Characteristics of *Myrtus communis* Linn.

Botanical Sources and Other Characteristics	General Description About This Plant	References
Botanical origin	Family: Myrtaceae Genus: -Myrtus Species name: <i>Myrtus communis</i> Linn (from Greece) Vernacular names (in Ethiopia): Ades (Amharic, Guragegna Tigregna), Addisaa, Coddoo (Affan Oromo); Wobattaa (Welaitigna)	1-4
Geographical origin	Only species of the genus found in the Northern Hemisphere, Southern Europe, North Africa, and West Asia Widespread in the Mediterranean region and others areas It is highly drought tolerant and needs only little to moderate water. It can grow in damp places, shades as well as full sun up to 800 m altitudes	1, 5, 6
Morphological characteristics of organized parts	Stem: Upright, 2.4-3 m tall, branched, thickly covered with evergreen leaves Leaves arrangement: Opposite, paired or whorled Leaves margin and size: Ovate to lanceolate with stiff structure, entire margined, acuminate, 2.5-3.8 cm long Flowers: Slender peduncles, medium size (2 cm diameter), stiff, has anthers Berries: Pea sized, orbicular	1, 5, 6
Organoleptic properties	Leaves: Dark green, glossy, glabrous, coriaceous Flowers: White, yellow anthers, fragrant smelling Berries: Blue black Fruits: Bitter when unripe, sweet when ripe Essential oil: Myrtle oil (very fragrant aromatic oil of this plant)	1, 2, 5, 6

and expanded by Arabs into an elaborate medical science based on the framework of the teaching of Hippocrates and Galen. It is a well-known shrub for its therapeutic, cosmetic, and food uses (Table 2). The name and use of *M communis* have been associated with myth and rituals in many societies.^{6,7}

Ethnopharmacological Studies

Extensive ethnopharmacological studies demonstrated that *M communis* L is endowed with promising pharmacological

**Figure 1.** *Myrtus communis* Linn.

activities, including antimicrobial, anticancer, antidiabetic, antiulcer, antidiarrheal, and anti-inflammatory activities.

Antimicrobial Effects

The leaves of *M communis* L revealed promising antimicrobial activities in the aqueous and ethanolic crude extracts,²⁵ fractionated constituents,^{26,27} a polar glycoside fraction, gallomyrtucommulone, isolated from the leaves,²⁸ in the biosynthesized silver nanoparticles.²⁹ The alcoholic extract of *M communis* was also found to be effective against selected extended-spectrum β -lactamase producing *Escherichia coli* isolates.³⁰ Also, the myrtle oil showed potent antimicrobial activities against *Helicobacter pylori*³¹ and clinical strains of *Mycobacterium tuberculosis*.³² It was further supported by the fact that α -pinene, 1,8 cineole, β -pinene, and limonene were found to be the highest contributors in antimicrobial properties of myrtle oil.³³ In situ results showed that the microbial population, except for *Pseudomonas* spp, decreased significantly in ground meat with added 5% of freeze-dried myrtle extract. The finding demonstrated the effectiveness of myrtle extract to control the proliferation of meat spoilage bacteria.³⁴ In a comparative study, patients treated with both *M communis* L and metronidazole base did not show any relapse; however, from those who were treated with metronidazole alone, 30% of them experienced relapsed during the follow-up period. From this it can be concluded that vaginal gel containing both the extract and the base is a more effective antibacterial than the base alone.³⁵ Besides, the isolated constituent, myrtacine, showed promising antibacterial activity against *Propionibacterium acnes* either alone or in combination with antibiotics and can be used for acne vulgaris treatment.³⁶

The ethnopharmacological studies of common myrtle extend to other microbial agents such as fungi and protozoa. All of the crude extracts of *M communis* have showed promising antifungal activity on *Candida albicans*.³⁷ What is more, myrtle oil was effective against all isolates of *Aspergillus* species.^{38,39} Another study revealed that the oil was also more active against *Cryptococcus neoformans* (yeast) and *Epidermophyton floccosum*, *Microsporium canis*, *Trichophyton rubrum* (dermatophytes).¹¹ Moreover, the oil inhibited growth of *Rhizoctonia solani* by 60% at the dose of 1600 ppm *in vitro*. Apart

Table 2. Ethnobotanical Uses of *Myrtus communis* Linn.

Parts Used	Description of Preparation	Application	Country (Used)	References
Leaves alone	Leaves have been used as an input for preparing liqueur called Mirto	Beverage	Italy	1, 4
	One gram of tea mixed with leaves has been drunk on daily basis	Stress and anxiety	Turkey	8
	Aqueous maceration of leaves can be taken after filtration and concentration	Depression, polymenorrhea, and wound	Iran	9
	The decoction of the leaf powder	Hypertension, eczema and other skin diseases	Algeria	10
	Not specified	Respiratory disorders and hemorrhoids		11
	Leaves are boiled and the stock is drunk (Turkey)	Abdominal pain and diarrhea	Pakistan, India, Turkey, Iran, Ethiopia	8, 11-14
	Juice of leaf is taken orally in the morning (Ethiopia), others unspecified			
	Rural women mix the leaf extract with raw butter and apply it to their hair	Cosmetics (hair fall control)	Ethiopia	4, 15
	Bathing with crushed fresh leaves	Dandruff	Ethiopia	14
	Dried leaf powder mixed with butter is applied topically	Scabies	Ethiopia	16
	The leaves are crushed, boiled with water, and are then drunk	Headache	Ethiopia	17
	The dried aqueous extract (leaf powder)	Sinus infections	China, France	18,19
	Dried leaf powder	Tinea capitis (“buha ras”) and as antipyretic and sedative agent	Ethiopia	20
Leaves, berries and myrtle oil	Depending on the nature of the disease states and parts of the plants used (remains unspecified)	Diarrhea, dysentery, gastric ulcer, vomiting, rheumatism, hemorrhages, deep sinuses, leucorrhea	India, Pakistan, Turkey, Ethiopia, Iran	6, 21, 22
		Hemorrhoid, inflammation, pulmonary and skin diseases		
		An astringent, antiseptic, disinfectant and hypoglycemic agent		
	The aqueous juice has also been used for the preparation of food and wines	Food and drinks	Europe (Italy)	6, 21
Leaves and fruits	Myrtle oil	Adjunct for the treatment of insomnia	Ethiopia	23
	Unspecified	Vulnerary, cough suppressant, and digestant effects	India, Pakistan	24

from this, both essential oil and methanolic extract of *M communis* leaf showed promising antileishmanial effect on *Leishmania tropica* in an *in vitro* model.⁴⁰ Myrtle oil also revealed antimalarial activity against *Plasmodium falciparum* *in vitro*.⁴¹

Anticancer Properties

Alwan et al⁴² showed that ethanolic extract of *M communis* leaves inhibited aryl hydrocarbon hydroxylase activity and 3H-benzo(a)pyrene binding to microsomal protein of rat liver, effectively. In the same study, no inhibitory effect was observed with aqueous extract.⁴² By the same authors of *in vitro* study, *M communis* showed significant inhibitory effect when *n*-butanol extract was used. The *n*-butanol extract was more effective than chloroform and petroleum ether extract. However, aqueous extract showed any inhibitory effects on both aryl hydrocarbon hydroxylase activity and 3H-benzo(a)pyrene binding to DNA *in vitro* unlike the *in vivo* study.⁴³ Myrtle oil also showed significant inhibition in cancer cell lines. At 200 µg/mL concentration, the inhibition was 67% and

95.2% for prostate and breast cancer cell lines, respectively. Probably the most exciting and hopeful result of this finding is that the inhibition value of essential oil of *M communis* on 3T3 fibroblast cell line is 3.7% and 6.5% for the dosages of 100 µg/mL and 200 µg/mL, respectively.⁴⁴ These results are so important and exciting since all of the aforementioned findings look like that sound of footsteps of the pioneer of an ideal and selective anticancer drug in the near future.⁴⁵

The cytotoxicity of myrtucommulone-A (MC-A) and its potential to induce apoptosis in cancer cells was demonstrated. MC-A showed an antiproliferative and strong inhibitor of migration of cancer cell. About 84 apoptotic pathway genes were screened to evaluate the effect of it on cancer cells. MC-A also mediated upregulation of apoptotic genes, including Fas, FasL, Gadd45a, Tnf, Tnfsf12, Trp53, and caspases through induction of both intrinsic and extrinsic apoptotic pathways.^{46,47} It can also induce apoptosis via activation of caspases (3, 8, and 9), cleavage of poly(ADP-ribose) polymerase (PARP), release of nucleosomes, and fragmentation of DNA. Moreover, it caused loss of the mitochondrial membrane

potential in MM6 cells and brought release of cytochrome *c* from mitochondria.⁴⁸ The activity of the myrtle oil was evaluated against spontaneous and *t*-BOOH (*t*-butyl hydroperoxide)–induced mutagenesis in *E coli* oxyR mutant IC202, a bacterial strain deficient in removing reactive oxygen species. When the oxidative mutagen was used, the oil expressed higher reduction of mutagenesis in a concentration-dependent manner.⁴⁹ Antimutagenic activity of myricetin-3-O-galactoside and myricetin-3-O-rhamnoside, isolated from the leaves of *M communis* L was assessed using the SOS chromotest and the Comet assay and they were found effective.⁵⁰

Antidiabetic Activity

The common myrtle demonstrated significant antihyperglycemic activity on its crude extracts against streptozotocin-induced diabetic mice.⁵¹ By the same token, the aqueous and methanolic extracts of *M communis* have significantly lowered blood glucose level in alloxan-induced diabetic mice. The aqueous extract has shown higher activity at relatively lower dose (500 mg/kg) than the methanolic extract at 1000 mg/kg dose. This indicates the aqueous extract is endowed with better efficacy compared with the methanolic extract as antidiabetic agent.⁵² A year later, another contradictory result, regarding aqueous and hydroalcoholic extracts, was also reported. Ethanolic extract of the leaves (2 g/kg) had a better hypoglycemic effect in diabetic rats compared with the aqueous extract ($P < .05$). Oral administration of the ethanolic extract (2 g/kg) had shown an additive effect on the hypoglycemic action of glibenclamide in rats.⁵³ Furthermore, data suggested that myrtle oil treatment reduces intestinal absorption of glucose, possibly via inhibiting α -glycosidase enzyme on alloxan-induced diabetic rabbits and on orally glucose loaded group.⁵⁴

Antiulcer Activity

Both methanol and aqueous crude extracts of the dried berries of myrtle revealed antiulcer activities against ethanol, indomethacin and pyloric ligation induced models in Wistar rats.⁵⁵ (–)-Myrtenol, isolated from the leaves of *M communis* L showed a significant reduction in the severity of ethanol-induced gastric lesions at all tested doses. The results provided an evidence for the gastroprotective effect of (–)-myrtenol that could be related to GABA_A-receptor activation and antioxidant activity.⁵⁶ A double-blind randomized controlled clinical trial revealed that *M communis* L freeze-dried aqueous extract showed comparable effect with omeprazole in gastrointestinal reflux disease.⁵⁷

Antidiarrheal Activity

In vitro studies on isolated tissue preparations demonstrated that hydroalcoholic (70% methanol) extract of *M communis* L possesses spasmolytic, bronchodilator, and vasodilator activities possibly due to blockade of voltage-dependent calcium channels and anticholinergic activity.⁵⁸ Furthermore, the




Figure 2. *Myrtus communis* L. as a candidate for the treatment of infectious diarrhea.

essential oil of the leaves of *M communis* possesses significant antidiarrheal and antispasmodic activity both *in vivo* and *in vitro*. The antidiarrheal activity produced by the oil was proved to be highly potent inhibitor of gastrointestinal motility and fluid secretion in mice and rat model of diarrhea, respectively. The oil also possesses antispasmodic activity in *ex vivo* model using isolated guinea pig ileum.⁵⁹ Similarly, an *in vivo* study found that myrtle berry seed aqueous extract administration induced a significant dose-dependent protection against diarrhea and intestinal fluid accumulation. The study revealed that myrtle berry seed aqueous extract had a potent protective effect against castor oil–induced acute diarrhea due in part to its antioxidant and antimicrobial properties.⁶⁰ Sisay et al⁶¹ also investigated the antidiarrheal activity of 80% methanol extract and solvent fraction of the leaves of *M communis* L in mice model of diarrhea *in vivo*. Therefore, it can be concluded that *M communis* L can be used for the treatment of diarrheas of diverse etiologies, including those with infectious component (Figure 2).

Antiinflammatory Activity

The 80% ethanol extract of *M communis* L leaf possessed anti-inflammatory activity on rats by measuring the suppression of carrageenan-induced paw edema using aspirin as the standard drug.⁶² It is further supported by isolated constituents (myrtucommulone, semimyrtucommulone, and nonprenylated acylphloroglucinols) that have also shown a promising anti-inflammatory activities.⁶³ MC-A isolated from myrtle showed the same pharmacological activity for inhibition of inflammation compared with the synthetic counterpart.⁴⁷ Treatment with *M communis* L ethanol extract reversed all the biochemical indices as well as histopathological alterations induced by acetic acid with the protective effects being similar to that of sulfasalazine treatment. The study showed that *M communis* L extract could alleviate colitis in rats and can be considered an alternative therapeutic approach for management of inflammatory bowel diseases.²¹ The anti-inflammatory potential of myrtle oil was also evaluated using an *in vitro* model of

Table 3. *Myrtus communis* L With Potential Antioxidant Activities.

Extracts	Methods Used	Result	Reference
Fruit crude extracts	1. DPPH and β -carotene-linoleic acid assays 2. Gas chromatography for fatty acid assay	High free radical scavenging activity in methanolic fruit extract	64
Chloroform, ethyl acetate, methanol, and aqueous extracts	1. DPPH assay 2. β -carotene bleaching assay 3. Ferric thiocyanate method 4. Thiobarbituric acid method	1. Ethyl acetate extract has the highest free radical scavenging power and reducing capability for DPPH and hydroxyl radical 2. Methanol extract exhibited higher chelating activity than ethyl acetate 3. Chloroform extract was strong inhibitor of lipid peroxidation in all assays	65
Leaves and berries Ethanol, methanol, water and ethyl acetate extracts of	1. Folin-Ciocalteu assay (total phenolic measurements) 2. Colorimetric method (total flavonoid measurements) 3. ABTS ⁺	1. Methanol and water extracts have significant antioxidant activities 2. Methanol > water > ethanol > ethyl acetate in both extracts (leaf and fruit) 3. Generally, the leaves have higher antioxidant activity than berries in phenolic and flavonoid content as well as activity	66
Berry extracts	1. HPLC coupled with electrospray mass spectrometry 2. Ultraviolet/visible detection 3. TEAC assay	Free radical scavenging activity. Antioxidant activity was preserved in 3 months	67
Myrtle oil	DPPH assay	The oil has moderate free radical scavenging activity	49
MBSAE extract	In isolated rat esophagus ER-induced damage	Potential protective and antioxidant effect	60

Abbreviations: DPPH, 2,2-diphenyl-1-picrylhydrazyl; ABTS⁺, 2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt; TEAC, trolox equivalent antioxidant capacity; ER, esophageal reflux; HPLC, high-performance liquid chromatography; MBSAE, myrtle berries seed aqueous extract.

lipopolysaccharide-stimulated macrophages. Assessment of cell viability was made through the MTT assay. The oil was able to significantly inhibit NO production, without affecting cell viability.¹¹

Antioxidant Activity

The antioxidant activity of various extracts of *M communis* L was investigated in extensive *in vitro* experimental methods as shown in Table 3.

Sedative-Hypnotic Activity

The essential oil of common myrtle revealed sedative-hypnotics.⁶⁸ Two years later, another controversial finding revealed that there was no induction of hypnosis. However, the myrtle oil prolonged pentobarbital-induced sleeping time and there was also half (50%) negative response on the chimney and traction test in a dose-dependent manner. The myrtle oil did not produce a hypnotic effect; however, it potentiated a hypnotic effect with significant central nervous system depressant activity.²³

Anxiolytic and Narcotic Analgesic Activity

Anxiolytic effect of 80% ethanolic extract of the leaves was investigated in rodent model. The data showed muscle relaxant

and anxiolytic effects of the extract with negligible anticonvulsant activity. The anxiolytic, myorelaxant, and hypnotic effects without effect on seizure threshold are in line with the effect of an $\alpha 2$ GABA receptor agonist.⁶⁹ The anxiolytic potential of myrtle oil was also investigated in rodent model.⁷⁰ These data were further supported by another finding in which a monoterpene alcohol, (–)myrtenol isolated from oil of myrtle presented anxiolytic-like activity that can be mediated by GABAergic transmission.⁷¹ Besides, its different crude extracts and isolated constituents showed narcotic analgesic properties.⁷²

Anthelmintic Activity

Findings of the study demonstrated that the myrtle oil at the concentration of 100 μ L/mL after 5 minutes of exposure killed 100% protoscolecies. Similarly, the mean mortality rate of protoscolecies after 10 minutes of exposure to concentration of 50 μ L/mL was 100%. The results showed potent scolicidal activity of *M communis* with no significant toxicity, which might be used as a natural scolicidal agent in hydatid cyst surgery.¹⁸

Antilipidemics and Antithrombotic Activity

The aqueous extract of *M communis* L leaves revealed antilipidemic and antithrombotic activities.⁷³




Figure 3. Five sesquiterpene-based meroterpenoids with 3 kinds of new skeletons [1, 2, 3, (+)-4, and (-)-4].

Pesticides

Myrtle oil has insect repellency effect, even with lower (10%) concentration.⁷⁴ Another finding suggested that the myrtle oil might have potential to be used as natural herbicides as well as fungicides.⁷⁵

Phytochemical Studies

The leaves of *M communis* were investigated to contain small amounts of phenolic acids (caffeic, ellagic, and gallic acids), and a flavonoid, quercetin derivatives (quercetin 3-O-galactoside and quercetin 3-O-rhamnoside). On the other hand, flavonoids such as galloyl derivatives of catechin and gallo catechin as well as myricetin derivatives are present in large amounts.⁷⁶ Similarly, 4 hydrolyzable tannins, 2 related polyphenolic compounds, and 4 myricetin glycosides were isolated from the leaves of *M communis* in another study.⁷⁷ The major terpenoids and their derivatives found in the essential oils of *M communis* L leaves were α -pinene, α -terpineol, linalool, 1,8-cineole, geranyl butyrate, geraniol, caryophyllene oxide, and neryl acetate.⁷⁸ Five sesquiterpene-based meroterpenoids with 3 kinds of new skeletons [1, 2, 3, (+)-4, and (-)-4] were isolated from the leaves of *M communis*⁷⁹ (Figure 3). It was further supported by both qualitative and quantitative studies. The chemical composition of 2 myrtle oil samples, taken from different localities, was investigated by gas chromatography–flame ionization detector, gas chromatography–mass spectrometry, and C^{13} nuclear magnetic resonance spectroscopy. Monoterpene derivatives were found to be the main compounds: α -pinene (50.8% and 33.6%), 1,8-cineole (21.9% and 13.3%), linalool (2.7% and 14.8%), and linalyl acetate (0.5% and 9.5%).¹¹ Chemical analysis of ethyl acetate extract revealed the presence of myricetin-3-O- α -rhamnoside. Ethyl acetate extract was found to have the highest total phenolic and total flavonoid contents with the values of 435.37 mg gallic acid equivalents/g dried weight and 130.75 mg quercetin equivalent/g dried weight, respectively.¹¹

Conclusion

The overall ethnobotanical, ethnopharmacological, and phytochemical studies of *M communis* L were well emphasized in this review article. Since time immemorial, this medicinal plant has played a significant role in the area of medicine and pharmacy as well as food and cosmetics applications. Majority of the folkloric reputes were validated by experimental studies, including antimicrobial, antidiarrheal, anticancer, antioxidant, antiulcer anti-inflammatory, and antidiabetic activities. Regarding the phytochemical analysis, sesquiterpene-based meroterpenoids were currently investigated in this plant and have shown to be associated with a multitude of biological activities.

Author Contributions

MS collected the literature, reviewed the literature critically, prepared the manuscript for publication, and read and approved the final version of the manuscript. TG edited the literature review and read and approved the final version of the manuscript.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

Ethical Approval

This article is a review study; therefore, no ethical approval is needed.

References

- Özkan AMG, Güray CG. A Mediterranean *Myrtus communis* L. (myrtle). In: Morel J-P, Mercuri Am, eds. *Plants and Culture: Seeds of the Cultural Heritage of Europe*. Bari, Italy: Centro Europeo per i Beni Culturali Ravello, Edipuglia Bari; 2009: 159-168.

2. Bouzabata A, Casanova J, Bighelli A, Cavaleiro C, Salgueiro L, Tomi F. The genus *Myrtus* L. in Algeria: composition and biological aspects of essential oils from *M. communis* and *M. nivellei*: a review. *Chem Biodivers*. 2016;13:672-680.
3. Hedberg I, Kelbessa E, Edwards S, et al. *Flora of Ethiopia and Eritrea*. 2006;5:539-540.
4. Tadesse M, Mesfin B. A review of selected plants used in the maintenance of health and wellness in Ethiopia. *Ethiopian e-J Res Innov Foresight*. 2010;2:85-102.
5. Aslam S, Ganaie KA, John AQ, et al. Family Myrtaceae in Kashmir *Myrtus communis* L: a new record for the shrub world of Kashmir Himalayas. *Acad Arena*. 2010;2:42-43.
6. Sumbul S, Ahmad MA, Asif M. *Myrtus communis* Linn: a review. *Indian J Nat Prod Resour*. 2011;2:395-402.
7. Husain A, Sofi GD, Tajuddin T, Dang R, Kumar N. Unani system of medicine—introduction and challenges. *Med J Islamic World Acad Sci*. 2010;18:27-30.
8. Akaydin G, Şimşek I, Arituluk ZC, et al. An ethno-botanical survey in selected towns of the Mediterranean sub region (Turkey). *Turk J Biol*. 2013;37:230-247.
9. Farzae MH, Abbasabadi Z, Shams-Ardekani MR. A comprehensive review of plants and their active constituents with wound healing activity in traditional Iranian medicine. *Wounds*. 2014; 26:197-206.
10. Sarri M, Mouyet FZ, Benziane M, Cheriet A. Traditional use of medicinal plants in a city at steppic character (M'sila, Algeria). *J Pharm Pharmacogn Res*. 2014;2:31-35.
11. Bouzabata A, Cabral C, Gonçalves MJ, Cruz MT, Bighelli A, Cavaleiro C. *Myrtus communis* L. as source of a bioactive and safe essential oil. *Food Chem Toxicol*. 2015;75:166-172.
12. Haq F, Ahmad H, Alam M. Traditional uses of medicinal plants of Nandiar Khuwarr catchment (District Battagram), Pakistan. *J Med Plants Res*. 2011;5:39-48.
13. Dogan Y, Ugulu I. Medicinal plants used for gastrointestinal disorders in some districts of Izmir Province, Turkey. *Ethnomedicine*. 2013;7:149-161.
14. Teklehaymanot T, Giday M. Ethno-botanical study of medicinal plants used in Zegie Peninsula, Northwestern Ethiopia. *J Ethnobiol Ethnomed*. 2007;3:12.
15. Tadesse M, Demissew S. Medicinal Ethiopian plants: inventory, identification and classification. In: Edwards S, Asfaw Z, eds. *Plants Used in African Traditional Medicine as Practiced in Ethiopia and Uganda, Botany 2000: East and Central Africa* (NAPRECA Monograph Series No. 5). Addis Ababa, Ethiopia: Addis Ababa University; 1992;1-24.
16. Gebeyehu G, Asfaw Z, Enyew A, et al. Ethno-botanical study of traditional medicinal plants and their conservation status in Mecha wereda, West Gojam Zone of Ethiopia. *Int J Pharm Health Care Res*. 2014;2:137-154.
17. Getaneh S, Girma Z. Ethnobotanical study of medicinal plants in Debrelibanose wereda, Central Ethiopia. *Afr J Plant Sci*. 2014;8: 366-379.
18. Mahmoudvand H, Fallahi S, Mahmoudvand H, Shakibaie M, Harandi MF, Dezaki ES. Efficacy of *Myrtus communis* L. to inactivate the hydatid cyst protoscoleces. *J Invest Surg*. 2016; 29:137-143.
19. Jabri MA, Rtibi K, Ben-Said A, et al. Antidiarrhoeal, antimicrobial and antioxidant effects of myrtle berries (*Myrtus communis* L.) seeds extract. *J Pharm Pharmacol*. 2016;68:264-274.
20. Jansen PCM. *Spices, Condiments and Medicinal Plants in Ethiopia: Their Taxonomy and Agricultural Significance*. Wageningen, the Netherlands: Centre for Agricultural Publishing and Documentation; 1981.
21. Sen A, Yuksel M, Bulut G, et al. Therapeutic potential of *Myrtus communis* subsp. *communis* extract against acetic acid-induced colonic inflammation in rats. *J Food Biochem*. 2016;41:e12297.
22. Alipour G, Dashti S, Hosseinzadeh H. Review of pharmacological effects of *Myrtus communis* L. and its active constituents. *Phytother Res*. 2014;28:1125-1136.
23. Birhanie MW, Walle B, Rebba K. Hypnotic effect of the essential oil from the leaves of *Myrtus communis* on mice. *Nat Sci Sleep*. 2016;8:267-275.
24. Tiwari S. Plants: a rich source of herbal medicine. *J Nat Prod*. 2008;1:27-35.
25. Ali MR, Hussin MS, Kadum MM, et al. In-vitro antimicrobial activities of *Myrtus communis* L. and *Cyperus rotundus* L. extracts. *Al- Mustansiriyah J Sci*. 2009;20:1.
26. Mansouri S, Alireza FA, Teimour GT, et al. Antibacterial activity of the crude extracts and fractionated constituents of *Myrtus communis*. *Pharm Biol*. 2001;39:399-401.
27. Alem G, Mekonnen Y, Tiruneh M, et al. In vitro antibacterial activity of crude preparation of myrtle (*Myrtus communis*) on common human pathogens. *Ethiopian Med J*. 2008;46:63-69.
28. Appendino G, Maxia L, Bettoni P, et al. Antibacterial galloylated alkylphloroglucinol glucosides from myrtle (*Myrtus communis*). *J Nat Prod*. 2006;69:251-254.
29. Sulaiman GM, Abd-ul-Wahed HE, Abd-ul-Wahab AI, et al. Antimicrobial activity of silver nanoparticles synthesized by *Myrtus communis* extract. *Eng Technol J*. 2013;31:3.
30. Saeidi S, Boroujeni NA, Ahmadi H, Hassanshahian M. Antibacterial activity of some plant extracts against extended-spectrum β -lactamase producing *Escherichia coli* isolates. *Jundishapur J Microbiol*. 2015;8:e15434.
31. Deriu A, Branca G, Molicotti P, et al. In vitro activity of essential oil of *Myrtus communis* L. against *Helicobacter pylori*. *Int J Antimicrob Agents*. 2007;30:562-563.
32. Zanetti S, Cannas S, Molicotti P, et al. Evaluation of the antimicrobial properties of the essential oil of *Myrtus communis* L. against clinical strains of *Mycobacterium* spp. *Interdiscip Perspect Infect Dis*. 2010;2010:931530.
33. Ebrahimabadi EH, Ghoreishi SM, Masoum S, Ebrahimabadi AH. Combination of GC/FID/mass spectrometry fingerprints and multivariate calibration techniques for recognition of antimicrobial constituents of *Myrtus communis* L. essential oil. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2016;1008:50-57.
34. Asaburi A., Di Martino V., Ercolini D., et al. Antimicrobial activity of *Myrtus communis* L. water-ethanol extract against meat spoilage strains of *Brochothrix thermosphacta* and *Pseudomonas fragi* in vitro and in meat. *Ann Microbiol*. 2015;65:841. doi:10. 1007/s13213-014-0925-9
35. Masoudi M, Miraj S, Kopaei MF. Comparison of the effects of *Myrtus communis* L, *Berberis vulgaris* and metronidazole vaginal

- gel alone for the treatment of bacterial vaginosis. *J Clin Diagn Res.* 2016;10:QC04-QC07.
36. Feuillolay C, Pecastaings S, Le Gac C, et al. A *Myrtus communis* extract enriched in myrtucummulones and ursolic acid reduces resistance of *Propionibacterium acnes* biofilms to antibiotics used in acne vulgaris. *Phytomedicine.* 2016;23:307-315.
 37. Erdogan EA, Goksen G, Everest A. Anti-candidal activities of some *Myrtus communis* L. extracts obtained using accelerated solvent extraction (ASE). *J Appl Biol Biotechnol.* 2014;2:12-14.
 38. Curini M, Bianchi A, Epifamo F, et al. In vitro antifungal activity of essential oils of *Erigeron canadensis* and *Myrtus communis* from France. *Chem Nat Comp.* 2003;30:191-194.
 39. Mohammadi R, Esfahani SH, Shadzi S, et al. Antifungal activity of *Myrtus communis* L essential oil against clinical isolates of *Aspergillus*. *J Isfahan Med Sch.* 2008;26:105-111.
 40. Mahmoudvand H, Ezzatkah F, Shariffar F. Antileishmanial and cytotoxic effects of essential oil and methanolic extract of *Myrtus communis* L. *Korean J Parasitol.* 2015;53:21-27.
 41. Milhau G, Valentin A, Benoit F, et al. In vitro antimalarial activity of 8 essential oils. *J Essential Oil Res.* 1997;9:329-333.
 42. Alwan AH, Al-Gaillany KA, Naji A. Inhibition of the binding of 3H-benzo(a)pyrene to rat liver microsomal protein by plant extracts. *Pharm Biol.* 1989;27:33-37.
 43. Alwan AH, Mahmoud MJ, Naji A. Effects of plant extracts on aryl hydrocarbon hydroxylase activity and 3H-benzo(a)pyrene binding to DNA. *J Food Saf.* 1989;10:209-218.
 44. Stevens N. Natural synergy: essential oils in cancer research. 2005. <http://www.youngliving.net/Presentations/StevensEssentialOilsInCancer.pdf>. Accessed August 1, 2014.
 45. Korkmaz A, Ogur R, Bakir B, Hasde M. Effects of AR-11 on human fibroblast, human lymphocytes, vero, hela, glioma, and MCF-7 cells Paper presented at: 2nd International Conference on Tumor Progression & Therapeutic Resistance; September 18-20, 2005; Boston, MA.
 46. Izgi K, Iskender B, Jauch J, et al. Myrtucummulone-A induces both extrinsic and intrinsic apoptotic pathways in cancer cells. *J Biochem Mol Toxicol.* 2015;29:432-439. doi:10.1002/jbt.21716.
 47. Hans M, Charpentier M, Huch V, et al. Stereoisomeric composition of natural myrtucummulone A. *J Nat Prod.* 2015;78:2381-2389.
 48. Tretiakova I, Blaesius D, Maxia L, et al. Myrtucummulone from *Myrtus communis* induces apoptosis in cancer cells via the mitochondrial pathway involving caspase-9. *Apoptosis.* 2008;13:119-131.
 49. Mimica-Dukic N, Bugarin D, Grbovic S, et al. Essential oil of *Myrtus communis* L as a potential antioxidant and antimutagenic agents. *Molecules.* 2010;15:2759-2770.
 50. Hayder N, Bouhlel I, Skandrani I, et al. In vitro antioxidant and antigenotoxic potentials of myricetin-3-O-galactoside and myricetin-3-O-rhamnoside from *Myrtus communis*: modulation of expression of genes involved in cell defence system using cDNA microarray. *Toxicol In Vitro.* 2008;22:567-581.
 51. Elfellah MS, Akhter MH, Khan MT. Anti-hyperglycaemic effect of an extract of *Myrtus communis* in streptozotocin-induced diabetes in mice. *J Ethnopharmacol.* 1984;11:275-281.
 52. Issa IA, Bule MH. A comparative study of the hypoglycemic effect of aqueous and methanolic extracts of *Myrtus communis* on alloxan induced diabetic Swiss albino mice. *Med Aromat Plants.* 2015;4:190. doi:10.4172/2167-0412.1000190.
 53. Panjeshahin MR, Azadbakht M, Akbari N. Antidiabetic activity of different extracts of *Myrtus communis* in streptozotocin induced diabetic rats. *Romanian J Diabetes Nutr Metab Dis.* 2016;23:183-190.
 54. Dineel AS, Acikgoz S, Cevik C, et al. Effects of in vivo antioxidant enzyme activities of myrtle oil in normoglycemic and alloxan diabetic rabbits. *J Ethnopharmacol.* 2007;110:498-503.
 55. Sumbul S, Ahmad MA, Asif M, et al. Evaluation of *Myrtus communis* L berries (common myrtle) in experimental ulcer models in rats. *Hum Exp Toxicol.* 2010;29:935-944.
 56. Viana AF, da Silva FV, Fernandes HD, Oliveira IS, Braga MA, Nunes PL. Gastroprotective effect of (–)-myrtenol against ethanol-induced acute gastric lesions: possible mechanisms. *J Pharm Pharmacol.* 2016;68:1085-1092.
 57. Zohalinezhad ME, Hosseini-Asl MK, Akrami R, et al. *Myrtus communis* L. freeze-dried aqueous extract versus omeprazol in gastrointestinal reflux disease: a double-blind randomized controlled clinical trial. *J Evid Based Complementary Altern Med.* 2016;21:23-29.
 58. Janbaz KH, Nisa M, Saqib F, et al. Bronchodilator, vasodilator and spasmolytic activities of methanolic extract of *Myrtus communis* L. *J Physiol Pharmacol.* 2013;64:479-484.
 59. Chala A, Engidawork E, Asres K. *Antidiarrheal and Antispasmodic Activities of Essential Oil of Myrtus communis L.* Addis Ababa, Ethiopia: School of Graduates, Addis Ababa University; 2011.
 60. Jabri M, Rtibi K, Said A, Aouadhi C, Hosni K, Sakly M. Anti-diarrhoeal, antimicrobial and antioxidant effects of myrtle berries (*Myrtus communis* L.) seeds extract. *J Pharm Pharmacol.* 2016;68:264-274.
 61. Sisay M, Engidawork E, Shibeshi W. Evaluation of the antidiarrheal activity of the leaf extracts of *Myrtus communis* Linn (Myrtaceae) in mice model. *BMC Complement Altern Med.* 2017;17:103. doi:10.1186/s12906-017-1625-3.
 62. Al-Hindawi MK, Al-Deen IH, Nabi MH, et al. Anti-inflammatory activity of some Iraqi plants using intact rats. *J Ethnopharmacol.* 1989;26:163-168.
 63. Feisst C, Franke L, Appendino G, Werz O. Identification of molecular targets of the oligomeric nonprenylated acylphloroglucinols from *Myrtus communis* and their implication as anti-inflammatory compounds. *J Pharmacol Exp Ther.* 2005;315:389-396.
 64. Serce S, Ercisli S, Sengul M, et al. Antioxidant activities and fatty acid composition of wild grown myrtle (*Myrtus communis* L.) fruits. *Pharmacogn Mag.* 2010;6:9.
 65. Bouaziz A, Khennouf S, Zarga MA, et al. Phytochemical analysis, hypotensive effect and antioxidant properties of *Myrtus communis* L. growing in Algeria. *Asian Pac J Trop Biomed.* 2015;5:19-28.
 66. Amensour M, Sendra E, Abrini J, et al. Antioxidant activity and total phenolic compounds of myrtle extracts. *J Food.* 2010;8:95-101.

67. Montoro P, Tuberoso CL, Piacente S, et al. Stability and antioxidant activity of polyphenols in extracts of *Myrtus communis* L. *J Pharm Biomed Anal.* 2006;41:1614-1619.
68. Walle M, Walle B, Zerihun L, et al. Sedative-hypnotic like effect of the essential oil from the leaves of *Myrtus communis* on mice. *Am J Biomed Life Sci.* 2014;2:70-77.
69. Hajiaghache R, Faizi M, Shahmohammadi Z, et al. Hydroalcoholic extract of *Myrtus communis* can alter anxiety and sleep parameters: a behavioural and EEG sleep pattern study in mice and rats. *Pharm Biol.* 2016;54:2141-2148. doi:10.3109/13880209.2016.1148175.
70. Hailu E, Engidawork E, Asres K. The essential oil of *Myrtus communis* L produces a non-sedating anxiolytic effect in mice models of anxiety. *Ethiopian Pharm J.* 2011;29:1-12.
71. Moreira MR, Salvadori MG, Almeida AA, et al. Anxiolytic-like effects and mechanism of (–)-myrtenol: a monoterpene alcohol. *Neurosci Lett.* 2014;579:119-124.
72. Twaij H, El-Jalil HA. Evaluation of narcotic (opioid like) analgesic activities of medicinal plants. *Eur J Sci Res.* 2009;33:179-182.
73. Khan RA, Feroz R, Jemil M, et al. Hypolipidemic and antithrombotic evaluation of aqueous extract of *Myrtus communis* L. *Afr J Pharm Pharmacol.* 2014;8:235-239.
74. Kayedi MH, Haghdoost AA, Salehnia A, et al. Evaluation of repellency effect of essential oils of *Satureja khuzestanica* (carvacrol), *Myrtus communis* (myrtle), *Lavendula officinalis* and *Salvia sclarea* using standard WHO repellency tests. *J Arthropod Borne Dis.* 2013;8:60-68.
75. Kordali S, Usanmaz A, Cakir A, Komaki A, Ercisli S. Antifungal and herbicidal effects of fruit essential oils of four *Myrtus communis* genotypes. *Chem Biodivers.* 2016;13:77-84.
76. Al-Hajjar MA, Al-Shahiry KF, Al-Abache SM. Isolation and identification of some active constituents of *Myrtus communis* in Iraq by using HPLC, FTIR techniques. *Iraqi Natl J Chem.* 2012;45:11-22.
77. Yoshimura M, Amakura Y, Tokuhara M, et al. Polyphenolic compounds isolated from the leaves of *Myrtus communis*. *J Nat Med.* 2008;62:366-368.
78. Khani A, Basavand F. Chemical composition and insecticidal activity of myrtle (*Myrtus communis* L.) essential oil against two stored-product pests. *J Med Plants Byprod.* 2012;2: 83-89.
79. Liu C, Ang S, Huang X. Meroterpenoids with new skeletons from *Myrtus communis* and structure revision of myrtucommulone K. *Org Lett.* 2016;18:4004-4007.